

# Lecture 2

## Presburger Arithmetic and Quantifier Elimination

Viktor Kuncak

# History of Presburger Arithmetic

Integer arithmetic with logical operations (and, or, not), quantifiers, only addition as an arithmetic operation, and  $<$  and  $=$  as a relation.

- ▶ minimalistically one can define a variant of it over non-negative natural numbers as having  $\wedge, \neg, \forall, +, =$  as the only symbols

One of the earliest theories shown decidable. Mojżesz Presburger gave an algorithm for quantifier elimination in 1929.

- ▶ a student of a famous logician Alfred Tarski
- ▶ Tarski gave him this question for his MSc thesis

The result at this time was of interest to mathematical logic and foundations of mathematics

- ▶ only much later it found applications in automated reasoning (Cooper 1972, Derek C. Oppen - STOC 1973)

# Presburger Arithmetic for Verification

res = 0

i = x

**while** // invariant  $I(res,i): res + 2*i == 2*x \ \&\& \ 0 \leq i$

(i > 0) {

  i = i - 1

  res = res + 2

}

Verification condition (VC) for preservation of loop invariant:

$$[I(res, i) \wedge i' = i - 1 \wedge res' = res + 2 \wedge 0 < i] \rightarrow I(res', i')$$

To prove that this VC is valid, we check whether its *negation*

$$I(res, i) \wedge i' = i - 1 \wedge res' = res + 2 \wedge 0 < i \wedge \neg I(res', i')$$

is *satisfiable*, i.e. whether this PA formula is true:

$$\begin{aligned} \exists x, res, i, res', i'. [ & res + 2i = 2x \wedge 0 \leq i \wedge 0 < i \wedge \\ & i' = i - 1 \wedge res' = res + 2 \wedge \\ & \neg(res' + 2i' = 2x \wedge 0 \leq i') ] \end{aligned}$$

## Introducing: One-Point Rule

If  $\bar{y}$  is a tuple of variables not containing  $x$ , then

$$\exists x.(x = t(\bar{y}) \wedge F(x, \bar{y})) \iff F(t(\bar{y}), \bar{y})$$

Proof:

- : Consider the values of  $\bar{y}$  such that there exists  $x$ , say  $x_1$ , for which  $x_1 = t(\bar{y}) \wedge F(x_1, \bar{y})$ . Because  $F(x_1, \bar{y})$  evaluates to true and the values of  $x_1$  and  $t(\bar{y})$  are the same,  $F(t, \bar{y})$  also evaluates to true.
- ← : Let  $\bar{y}$  be such that  $F(t, \bar{y})$  holds. Let  $x$  be the value of  $t(\bar{y})$ . Then of course  $x = t(\bar{y})$  evaluates to true and so does  $F(x, \bar{y})$ . So there exists  $x$  for which  $x = t(\bar{y}) \wedge F(x, \bar{y})$  holds.

*One point rule:*

replaces left side (LHS) of equivalence by the right side (RHS).

*Flattening*, used when  $t$  is complex, replaces RHS by LHS.

## Dual One-Point Rule for $\forall$

$$\forall x.(x = t(\bar{y}) \rightarrow F(x, \bar{y})) \iff F(t(\bar{y}), \bar{y})$$

To prove it, negate both sides:

$$\exists x.(x = t(\bar{y}) \wedge \neg F(x, \bar{y})) \iff \neg F(t(\bar{y}), \bar{y})$$

so it reduces to the rule for  $\exists$ .

## Using One-Point Rule on Negated Verification Condition

$$\exists x, res, i, res', \underline{i}'. \left[ res + 2i = 2x \wedge 0 \leq i \wedge 0 < i \wedge \right. \\ \left. \underline{i}' = i - 1 \wedge res' = res + 2 \wedge \right. \\ \left. \neg(res' + 2i' = 2x \wedge 0 \leq i') \right]$$

## Using One-Point Rule on Negated Verification Condition

$$\exists x, res, i, res', \underline{i}'. \left[ res + 2i = 2x \wedge 0 \leq i \wedge 0 < i \wedge \right. \\ \left. \underline{i' = i - 1} \wedge res' = res + 2 \wedge \right. \\ \left. \neg(res' + 2i' = 2x \wedge 0 \leq i') \right]$$

$$\exists x, res, i, \underline{res}'. \left[ res + 2i = 2x \wedge 0 \leq i \wedge 0 < i \wedge \right. \\ \left. \underline{res' = res + 2} \wedge \right. \\ \left. \neg(res' + 2(i - 1) = 2x \wedge 0 \leq i - 1) \right]$$

## Using One-Point Rule on Negated Verification Condition

$$\exists x, res, i, res', \underline{i}'. \left[ res + 2i = 2x \wedge 0 \leq i \wedge 0 < i \wedge \right. \\ \left. \underline{i' = i - 1} \wedge res' = res + 2 \wedge \right. \\ \left. \neg(res' + 2i' = 2x \wedge 0 \leq i') \right]$$

$$\exists x, res, i, \underline{res}'. \left[ res + 2i = 2x \wedge 0 \leq i \wedge 0 < i \wedge \right. \\ \left. \underline{res' = res + 2} \wedge \right. \\ \left. \neg(res' + 2(i - 1) = 2x \wedge 0 \leq i - 1) \right]$$

$$\exists x, res, i. \left[ \underline{res + 2i = 2x} \wedge 0 \leq i \wedge 0 < i \wedge \right. \\ \left. \neg(res + 2 + 2(i - 1) = 2x \wedge 0 \leq i - 1) \right]$$

## Using One-Point Rule on Negated Verification Condition

$$\exists x, res, i, res', \underline{i}'. \left[ res + 2i = 2x \wedge 0 \leq i \wedge 0 < i \wedge \right. \\ \left. \underline{i' = i - 1} \wedge res' = res + 2 \wedge \right. \\ \left. \neg(res' + 2i' = 2x \wedge 0 \leq i') \right]$$

$$\exists x, res, i, \underline{res}'. \left[ res + 2i = 2x \wedge 0 \leq i \wedge 0 < i \wedge \right. \\ \left. \underline{res' = res + 2} \wedge \right. \\ \left. \neg(res' + 2(i - 1) = 2x \wedge 0 \leq i - 1) \right]$$

$$\exists x, res, i. \left[ \underline{res + 2i = 2x} \wedge 0 \leq i \wedge 0 < i \wedge \right. \\ \left. \neg(res + 2 + 2(i - 1) = 2x \wedge 0 \leq i - 1) \right]$$

$$\exists x, \underline{res}, i. \left[ \underline{res = 2x - 2i} \wedge 0 \leq i \wedge 0 < i \wedge \right. \\ \left. \neg(res + 2 + 2(i - 1) = 2x \wedge 0 \leq i - 1) \right]$$

## Using One-Point Rule on Negated Verification Condition

$$\exists x, res, i, res', \underline{i}'. \left[ \begin{aligned} &res + 2i = 2x \wedge 0 \leq i \wedge 0 < i \wedge \\ &\underline{i' = i - 1} \wedge res' = res + 2 \wedge \\ &\neg(res' + 2i' = 2x \wedge 0 \leq i') \end{aligned} \right]$$

$$\exists x, res, i, \underline{res}'. \left[ \begin{aligned} &res + 2i = 2x \wedge 0 \leq i \wedge 0 < i \wedge \\ &\underline{res' = res + 2} \wedge \\ &\neg(res' + 2(i - 1) = 2x \wedge 0 \leq i - 1) \end{aligned} \right]$$

$$\exists x, res, i. \left[ \begin{aligned} &\underline{res + 2i = 2x} \wedge 0 \leq i \wedge 0 < i \wedge \\ &\neg(res + 2 + 2(i - 1) = 2x \wedge 0 \leq i - 1) \end{aligned} \right]$$

$$\exists x, \underline{res}, i. \left[ \begin{aligned} &\underline{res = 2x - 2i} \wedge 0 \leq i \wedge 0 < i \wedge \\ &\neg(res + 2 + 2(i - 1) = 2x \wedge 0 \leq i - 1) \end{aligned} \right]$$

$$\exists x, i. \left[ \begin{aligned} &0 \leq i \wedge 0 < i \wedge \\ &\neg(2x - 2i + 2 + 2(i - 1) = 2x \wedge 0 \leq i - 1) \end{aligned} \right]$$

Simplifies to  $\exists x, i. 0 < i \wedge \neg(0 \leq i - 1)$  and then to false.

$$\begin{aligned} &1 \leq i \\ &\neg(1 \leq i) \end{aligned}$$

But there is more

One-point rule is one of the many steps used in **quantifier elimination** procedures.

## Quantifier Elimination (QE)



Given a formula  $F(\bar{y})$  containing quantifiers find a formula  $G(\bar{y})$

- ▶ **equivalent** to  $F(\bar{y})$
- ▶ that has **no quantifiers**
- ▶ and has a **subset (or equal set) of free variables** of  $F$

Note

- ▶ Equivalence: For all  $\bar{y}$ ,  $F(\bar{y})$  and  $G(\bar{y})$  have same truth value  
 $\leadsto$  we can use  $G(\bar{y})$  instead of  $F(\bar{y})$
- ▶ No quantifiers: easier to check satisfiability of  $G(\bar{y})$

$\bar{y}$  is a possibly empty tuple of variables

## We are lucky when a theory has (“admits”) QE

Suppose  $F$  has no free variables (all variables are quantified).  
What is the result of applying QE to  $F$ ?

## We are lucky when a theory has (“admits”) QE

Suppose  $F$  has no free variables (all variables are quantified).

What is the result of applying QE to  $F$ ?

Are there any variables in the resulting formula?

## We are lucky when a theory has (“admits”) QE

Suppose  $F$  has no free variables (all variables are quantified).

What is the result of applying QE to  $F$ ?

Are there any variables in the resulting formula?

- ▶ No free variables: they are a subset of the original, empty set
- ▶ No quantified variables: because it has no quantifiers 😊

Formula without any variables! Example:

$$(2 + 4 = 7) \vee (1 + 1 = 2)$$

We check the truth value of such formula by simply evaluating it!

## Using QE for Deciding Satisfiability/Validity

- ▶ To check satisfiability of  $H(\bar{y})$ : eliminate the quantifiers from  $\exists \bar{y}.H(\bar{y})$  and evaluate.
- ▶ Validity: eliminate quantifiers from  $\forall \bar{y}.H(\bar{y})$  and evaluate

We can even check formulas like this:

$$\forall x, y, r. \exists z. (5 \leq r \wedge x + r \leq y) \rightarrow (x < z \wedge z < y \wedge 3|z)$$

Here  $3|z$  denotes that  $z$  is divisible by 3.

Does Presburger Arithmetic admit QE?

# Does Presburger Arithmetic admit QE?

Depends on the particular set of symbols!

Given a formula  $F(\bar{y})$  containing quantifiers find a formula  $G(\bar{y})$

- ▶ **equivalent** to  $F(\bar{y})$
- ▶ that has **no quantifiers**
- ▶ and has a **subset (or equal set) of free variables** of  $F$

If we lack some operations that can be expressed using quantifiers, there may be no equivalent formula without quantifiers.

- ▶  $\exists y.x = y + y + y$ , so we better have divisibility

Quantifier elimination says: if you can define some relationship between variables using an arbitrary, possibly quantified, formula  $F$ ,

$$r \stackrel{\text{def}}{=} \{(x, y) \mid F(x, y)\}$$

then you can also define same  $r$  using another quantifier-free formula  $G$ .

# Presburger Arithmetic (PA)

We look at the theory of integers with addition.

- ▶ introduce constant for each integer constant
- ▶ to be able to restrict values to natural numbers when needed, and to compare them, we introduce  $<$
- ▶ introduce not only addition but also subtraction
- ▶ to conveniently express certain expressions, introduce function  $m_K$  for each  $K \in \mathbb{Z}$ , to be interpreted as multiplication by a constant,  $m_K(x) = K \cdot x$ . We write  $m_K$  as  $K \cdot x$ .  
Note: there is *no multiplication between variables* in PA
- ▶ to enable quantifier elimination from  $\exists x.y = K \cdot x$  introduce for each  $K$  predicate  $K|y$  (divisibility,  $y \% K = 0$ )

The resulting language has these function and relation symbols:

$\{+, -, =, <\} \cup \{K \mid K \in \mathbb{Z}\} \cup \{(K \cdot -) \mid K \in \mathbb{Z}\} \cup \{(K| -) \mid K \in \mathbb{Z}\}$

We also have, as usual:  $\wedge, \vee, \neg, \rightarrow$  and also:  $\exists, \forall$

## Example

**Eliminate  $y$  from this formula:**

$$\exists y. (3y - 2w + 1 > -w \wedge 2y - 6 < z \wedge 4 \mid 5y + 1)$$

What should we do first?

## Example

**Eliminate  $y$  from this formula:**

$$\exists y. 3y - 2w + 1 > -w \wedge 2y - 6 < z \wedge 4 \mid 5y + 1$$

What should we do first?

Simplify/normalize what we can using properties of integer operations:

$$\exists y. 0 < -w + 3y + 1 \wedge 0 < -2y + z + 6 \wedge 4 \mid 5y + 1$$

## Example

**Eliminate  $y$  from this formula:**

$$\exists y. 3y - 2w + 1 > -w \wedge 2y - 6 < z \wedge 4 \mid 5y + 1$$

What should we do first?

Simplify/normalize what we can using properties of integer operations:

$$\exists y. 0 < -w + 3y + 1 \wedge 0 < -2y + z + 6 \wedge 4 \mid 5y + 1$$

First we will consider only eliminating existential from a **conjunction of literals**.

# Conjunctions of Literals

**Atomic formula:** a relation applied to argument.

Here, relations are:  $=$ ,  $<$ ,  $K|_-$ . So, atomic formulas are:

$$t_1 = t_2, \quad t_1 < t_2, \quad K | t$$

# Conjunctions of Literals

**Atomic formula:** a relation applied to argument.

Here, relations are:  $=$ ,  $<$ ,  $K|_-$ . So, atomic formulas are:

$$t_1 = t_2, \quad t_1 < t_2, \quad K | t$$

**Literal:** Atomic formula or its negation. Example:  $\neg(x = y + 1)$

Conjunction of literals:  $L_1 \wedge \dots \wedge L_n$

- ▶ no disjunctions, no implications
- ▶ negation only applies to atomic formulas

We first consider the quantifier elimination problem of the form:

$$\exists y. L_1 \wedge \dots \wedge L_n$$

This will prove to be sufficient to eliminate all quantifiers!

## Eliminating $\exists$ from conjunction of literals suffices

Can we eliminate  $\exists$  from any **quantifier-free formula**?

$$\exists x.F(x, \bar{y})$$

where  $F$  is quantifier-free?

## Eliminating $\exists$ from conjunction of literals suffices

Can we eliminate  $\exists$  from any **quantifier-free formula**?

$$\exists x.F(x, \bar{y})$$

where  $F$  is quantifier-free?

Formula without quantifiers has  $\wedge, \vee, \neg$  applied to atomic formulas.

## Eliminating $\exists$ from conjunction of literals suffices

Can we eliminate  $\exists$  from any **quantifier-free formula**?

$$\exists x.F(x, \bar{y})$$

where  $F$  is quantifier-free?

Formula without quantifiers has  $\wedge, \vee, \neg$  applied to atomic formulas.

Convert  $F$  to **disjunctive normal form**:

$$F \iff \bigvee_{i=1}^m C_i$$

each  $C_i$  is a **conjunction of literals**.

## Eliminating $\exists$ from conjunction of literals suffices

Can we eliminate  $\exists$  from any **quantifier-free formula**?

$$\exists x.F(x, \bar{y})$$

where  $F$  is quantifier-free?

Formula without quantifiers has  $\wedge, \vee, \neg$  applied to atomic formulas.

Convert  $F$  to **disjunctive normal form**:

$$F \iff \bigvee_{i=1}^m C_i$$

each  $C_i$  is a **conjunction of literals**.

$$\left[ \exists x. \bigvee_{i=1}^m C_i \right] \iff \bigvee_{i=1}^m (\exists x. C_i)$$

# How does disjunctive normal form (DNF) transformation work?

Which steps should we use?

# How does disjunctive normal form (DNF) transformation work?

Which steps should we use?

**Negation propagation:**

$$\neg(p \wedge q) \rightsquigarrow (\neg p) \vee (\neg q)$$

$$\neg(p \vee q) \rightsquigarrow (\neg p) \wedge (\neg q)$$

$$\neg\neg p \rightsquigarrow p$$

Result is **negation-normal form**, NNF

NNF transformation is polynomial (exercise!)

# How does disjunctive normal form (DNF) transformation work?

Which steps should we use?

**Negation propagation:**

$$\neg(p \wedge q) \rightsquigarrow (\neg p) \vee (\neg q)$$

$$\neg(p \vee q) \rightsquigarrow (\neg p) \wedge (\neg q)$$

$$\neg\neg p \rightsquigarrow p$$

Result is **negation-normal form**, NNF

NNF transformation is polynomial (exercise!)

**Distributivity**

$$a \wedge (b_1 \vee b_2) \rightsquigarrow (a \wedge b_1) \vee (a \wedge b_2)$$

This can lead to exponential explosion.

Can we obtain equivalent DNF formula without explosion?

# How does disjunctive normal form (DNF) transformation work?

Which steps should we use?

**Negation propagation:**

$$\neg(p \wedge q) \rightsquigarrow (\neg p) \vee (\neg q)$$

$$\neg(p \vee q) \rightsquigarrow (\neg p) \wedge (\neg q)$$

$$\neg\neg p \rightsquigarrow p$$

Result is **negation-normal form**, NNF

NNF transformation is polynomial (exercise!)

**Distributivity**

$$a \wedge (b_1 \vee b_2) \rightsquigarrow (a \wedge b_1) \vee (a \wedge b_2)$$

This can lead to exponential explosion.

Can we obtain equivalent DNF formula without explosion?

No! See exercise.

## Eliminating from quantifier free formulas

$$\exists x.F \iff \left[ \exists x. \bigvee_{i=1}^m C_i \right] \iff \bigvee_{i=1}^m (\exists x.C_i)$$

## Nested Existential Quantifiers

$$\exists x_1. \exists x_2. \exists x_3. F_0(x_1, x_2, x_3, \bar{y})$$

## Nested Existential Quantifiers

$$\exists x_1. \exists x_2. \exists x_3. F_0(x_1, x_2, x_3, \bar{y})$$

$$\exists x_1. \exists x_2. F_1(x_1, x_2, \bar{y})$$

## Nested Existential Quantifiers

$$\exists x_1. \exists x_2. \exists x_3. F_0(x_1, x_2, x_3, \bar{y})$$

$$\exists x_1. \exists x_2. F_1(x_1, x_2, \bar{y})$$

$$\exists x_1. F_2(x_1, \bar{y})$$

# Nested Existential Quantifiers

$$\exists x_1. \exists x_2. \exists x_3. F_0(x_1, x_2, x_3, \bar{y})$$

$$\exists x_1. \exists x_2. F_1(x_1, x_2, \bar{y})$$

$$\exists x_1. F_2(x_1, \bar{y})$$

$$F_3(\bar{y})$$

# Nested Existential Quantifiers

$$\exists x_1. \exists x_2. \exists x_3. F_0(x_1, x_2, x_3, \bar{y})$$

$$\exists x_1. \exists x_2. F_1(x_1, x_2, \bar{y})$$

$$\exists x_1. F_2(x_1, \bar{y})$$

$$F_3(\bar{y})$$



# Universal Quantifiers

If  $F_0(x, \bar{y})$  is quantifier-free, how to eliminate

$$\forall y. F_0(x, \bar{y})$$

## Universal Quantifiers

If  $F_0(x, \bar{y})$  is quantifier-free, how to eliminate

$$\forall y. F_0(x, \bar{y})$$

Equivalence (property always holds if there is no counterexample):

$$\forall y. F_0(x, \bar{y}) \iff \neg[\exists y. \neg F_0(x, \bar{y})]$$

It thus suffices to process:

$$\neg[\exists y. \neg F_0(x, \bar{y})]$$

Note that  $\neg F_0(x, \bar{y})$  is quantifier-free, so we know how to handle it:

$$\exists y. \neg F_0(x, \bar{y}) \rightsquigarrow F_1(\bar{y})$$

Therefore, we obtain

$$\neg F_1(\bar{y})$$

## Removing any **alternation** of quantifiers: illustration

Alternation: switch between existentials and universals

$$\exists x_1. \forall x_2. \forall x_3. \exists x_4. F_0(x_1, x_2, x_3, x_4, \bar{y})$$

$$\exists x_1. \neg \exists x_2. \exists x_3. \neg \exists x_4. \underline{F_0(x_1, x_2, x_3, x_4, \bar{y})}$$

$$\exists x_1. \neg \exists x_2. \underline{\exists x_3. \neg F_1(x_1, x_2, x_3, \bar{y})}$$

$$\exists x_1. \neg \exists x_2. \underline{F_2(x_1, x_2, \bar{y})}$$

$$\exists x_1. \neg F_3(x_1, \bar{y})$$

$$F_4(\bar{y})$$

Each quantifier alternation involves a disjunctive normal form transformation.

In practice, we do not have many alternations.

# Back to Presburger Arithmetic

Consider the quantifier elimination problem of the form:

$$\exists y. L_1 \wedge \dots \wedge L_n$$

where  $L_i$  are literals from PA.

Note that, for integers:

- ▶  $\neg(x < y) \iff y \leq x$
- ▶  $x < y \iff x + 1 \leq y$
- ▶  $x \leq y \iff x < y + 1$

We use these observations below.

Instead of  $\leq$  we choose to use  $<$  only.

We do not write  $x > y$  but only  $y < x$ .

# Normalizing Literals for PA

**Normal Form of Terms:** All *terms* are built from  $K, +, -, K \cdot \neg$ , so using standard transformations they can be represented as:

$K_0 + \sum_{i=1}^n K_i x_i$  We call such term a linear term.

**Normal Form for Literals in PA:**

$$\neg(t_1 < t_2) \text{ becomes } t_2 < t_1 + 1$$
$$\neg(t_1 = t_2) \text{ becomes } t_1 < t_2 \vee t_2 < t_1$$

*Handwritten notes:*  $\exists y. x < y \wedge \neg (\exists |x)$   
 $\exists y. [x < y \wedge (\exists |x+1 \vee \exists |x+2)]$   
 $\exists y. [(x < y \wedge \exists |x+1) \vee (x < y \wedge \exists |x+2)]$

$$t_1 = t_2 \text{ becomes } t_1 < t_2 + 1 \wedge t_2 < t_1 + 1 \quad (*)$$

$$\neg(K | t) \text{ becomes } \bigvee_{i=1}^{K-1} K | t + i$$

$$t_1 < t_2 \text{ becomes } 0 < t_2 - t_1$$

To remove disjunctions we generated, compute DNF again.

(\*) We transformed equalities just for simplicity. Usually we handle them directly.

# Why one-point rule will not be enough

Need to handle inequalities, not just equalities

If we have integers, we cannot always divide perfectly.

Variable to eliminate can occur not as  $y$  but as, e.g.  $3y$

## Exposing the Variable to Eliminate: Example

$$\begin{array}{l} k | t \\ \exists i. t = i \cdot k \\ \exists i. c \cdot t = c \cdot i \cdot k \\ c \cdot t | c \cdot k \end{array}$$

$$\exists y. 0 < -w + \underline{3y} + 1 \wedge 0 < -\underline{2y} + z + 6 \wedge 4 | \underline{5y} + 1$$

Least common multiple of coefficients next to  $y$ ,

$$M = \text{lcm}(3, 2, 5) = 30$$

Make all occurrences of  $y$  in the body have this coefficient:

$$\exists y. 0 < -10w + \underline{30y} + 10 \wedge 0 < -\underline{30y} + 15z + 90 \wedge 24 | \underline{30y} + 6$$

Now we are quantifying over  $y$  and using  $30y$  everywhere.

Let  $x$  denote  $30y$ .

It is **not an arbitrary**  $x$ . It is divisible by 30.

$$\exists x. 0 < -10w + x + 10 \wedge 0 < -x + 15z + 90 \wedge 24 | x + 6 \wedge 30 | x$$

## Exposing the Variable to Eliminate in General

Eliminating  $y$  from conjunction  $F(y)$  of literals:

- ▶  $0 < t$
- ▶  $K \mid t$

where  $t$  is a linear term. To eliminate  $\exists y$  from such conjunction, we wish to ensure that the coefficient next to  $y$  is one or minus one.

Observation:

- ▶  $0 < t$  is equivalent to  $0 < c t$
- ▶  $K \mid t$  is equivalent to  $c K \mid c t$

for  $c$  a positive integer.

Let  $K_1, \dots, K_n$  be all coefficients next to  $y$  in the formula.

Let  $M$  be a positive integer such that  $K_i \mid M$  for all  $i$ ,  $1 \leq i \leq n$

- ▶ for example, let  $M$  be the **least common multiple**

$$M = \text{lcm}(K_1, \dots, K_n)$$

## Ensuring Coefficient One

Multiply each literal where  $y$  occurs in subterm  $K_i y$  by constant  $M/|K_i|$

- ▶ the point is,  $M$  is divisible by  $|K_i|$  by construction

What is the coefficient next to  $y$  in the resulting formula?

## Ensuring Coefficient One

Multiply each literal where  $y$  occurs in subterm  $K_i y$  by constant  $M/|K_i|$

- ▶ the point is,  $M$  is divisible by  $|K_i|$  by construction

What is the coefficient next to  $y$  in the resulting formula?

$M$  or  $-M$

## Ensuring Coefficient One

Multiply each literal where  $y$  occurs in subterm  $K_i y$  by constant  $M/|K_i|$

- ▶ the point is,  $M$  is divisible by  $|K_i|$  by construction

What is the coefficient next to  $y$  in the resulting formula?

$$M \text{ or } -M$$

We obtain a formula of the form  $\exists y. F(M \cdot y)$ .

Letting  $x = My$ , we conclude the formula is equivalent to

$$\exists x. F(x) \wedge (M \mid x)$$

What is the coefficient next to  $y$  in the resulting formula?

## Ensuring Coefficient One

Multiply each literal where  $y$  occurs in subterm  $K_i y$  by constant  $M/|K_i|$

- ▶ the point is,  $M$  is divisible by  $|K_i|$  by construction

What is the coefficient next to  $y$  in the resulting formula?

$$M \text{ or } -M$$

We obtain a formula of the form  $\exists y. F(M \cdot y)$ .

Letting  $x = My$ , we conclude the formula is equivalent to

$$\exists x. F(x) \wedge (M \mid x)$$

What is the coefficient next to  $y$  in the resulting formula?

$$1 \text{ or } -1$$

Lower and upper bounds:

$$\exists x. [F(x, y) \wedge G(y)] \\ (\exists x. [F(x, y)]) \wedge G(y)$$

Consider the coefficient next to  $x$  in  $0 < t$ . If it is  $-1$ , move the term to left side. If it is  $1$ , move the remaining terms to the left side. We obtain formula  $F_1(x)$  of the form

$$\bigwedge_{i=1}^L a_i + 1 \leq x \wedge \bigwedge_{j=1}^U x \leq b_j - 1 \wedge \bigwedge_{i=1}^D K_i \mid (x + t_i)$$

$$\exists x. a < x \wedge x < b$$

If there are no divisibility constraints ( $D = 0$ ), what is the formula equivalent to?

## Lower and upper bounds:

Consider the coefficient next to  $x$  in  $0 < t$ . If it is  $-1$ , move the term to left side. If it is  $1$ , move the remaining terms to the left side. We obtain formula  $F_1(x)$  of the form

$$\underbrace{\bigwedge_{i=1}^L a_i < x \wedge \bigwedge_{j=1}^U x < b_j \wedge \bigwedge_{i=1}^D K_i \mid (x + t_i)}_{F_1(x)}$$

If there are no divisibility constraints ( $D = 0$ ), what is the formula equivalent to?

$$\overbrace{\max_i a_i}^x + 1 \leq \min_j b_j - 1 \text{ which is equivalent to } \bigwedge_{ij} a_i + 1 < b_j$$

## Replacing variable by test terms

There is an alternative way to express the above condition by replacing  $F_1(x)$  with  $\bigvee_k F_1(t_k)$  where  $t_k$  do not contain  $x$ . This is a common technique in quantifier elimination. Note that if  $F_1(t_k)$  holds then certainly  $\exists x.F_1(x)$ .

What are example terms  $t_i$  when  $D = 0$  and  $L > 0$ ? Hint: ensure that at least one of them evaluates to  $\max a_j + 1$ .

$$\exists x.F_1(x) \quad \curvearrowright \quad \bigvee_{k=1}^L F_1(a_k + 1)$$

What if  $D > 0$  i.e. we have additional divisibility constraints?

$$\bigvee_{k=1}^L \bigvee_{i=1}^N F_1(a_k + i)$$

What is  $N$ ? least common multiple of  $K_1, \dots, K_D$

Note that if  $F_1(u)$  holds then also  $F_1(u - N)$  holds.

## Back to Example

$$\exists x. -10 + 10w < x \wedge x < 90 + 15z \wedge 24 \mid x + 6 \wedge 30 \mid x$$

## Back to Example

$$\exists x. -10 + 10w < x \wedge x < 90 + 15z \wedge 24 \mid x + 6 \wedge \underline{30 \mid x}$$

$$-10 + 10w < -10 + 10w + i \quad \underline{-10 + 10w + i} < \underline{90 + 15z}$$

$$\bigvee_{i=1}^{120} 10w + i < \underline{100} + 15z \wedge 0 < i \wedge 24 \mid 10w - 4 + i \wedge \underline{30 \mid 10w - 10 + i}$$

## Special cases

What if  $L = 0$ ? We first drop all constraints except divisibility, obtaining  $F_2(x)$

$$\bigwedge_{i=1}^D K_i \mid (x + t_i)$$

and then eliminate quantifier as

$$\bigvee_{i=1}^N F_2(i)$$

## It works

We finished describing a complete quantifier elimination algorithm for Presburger Arithmetic!

## It works

We finished describing a complete quantifier elimination algorithm for Presburger Arithmetic!

This algorithm and its correctness prove that:

- ▶ PA admits quantifier elimination
- ▶ Satisfiability, validity, entailment, equivalence of PA formulas is decidable

We can use the algorithm to prove verification conditions.

Even if not the most efficient way, it gives us insights on which we can later build to come up with better algorithms.

- ▶ Quantified and quantifier-free formulas have the same expressive power

Many other properties follow (e.g. interpolation).

# Interpolation For Logical Theories

$$x \geq 0 \vDash -10 \leq x$$

Interpolation can be useful in generalizing counterexamples to invariants.

Universal **Entailment**: we will write  $F_1 \models F_2$  to denote that for all free variables of  $F_1$  and  $F_2$ , if  $F_1$  holds then  $F_2$  holds.

Given two formulas such that

$$F_0(\bar{x}, \bar{y}) \models F_1(\bar{y}, \bar{z})$$

an interpolant for  $F_0, F_1$  is a formula  $I(\bar{y})$ , which has only variables common to  $F_0$  and  $F_1$ , such that

- ▶  $F_0(\bar{x}, \bar{y}) \models I(\bar{y})$ , and
- ▶  $I(\bar{y}) \models F_1(\bar{y}, \bar{z})$

In other words, the entailment between  $F_0$  and  $F_1$  can be explained through  $I(\bar{y})$ .

Logic has **interpolation property** if, whenever  $F_0 \models F_1$ , then there exists an interpolant for  $F_0, F_1$ .

We often wish to have *simple* interpolants, for example ones that are quantifier free.

# Quantifier Elimination Implies Interpolation

If logic has QE, it also has quantifier-free interpolants.

Consider the formula

$$\forall \bar{x}, \bar{y}, \bar{z}. [F_0(\bar{x}, \bar{y}) \rightarrow F_1(\bar{y}, \bar{z})]$$

$$\forall x. (p(x) \rightarrow q)$$

$$\forall x. (\neg p(x) \vee q)$$

pushing  $\bar{x}$  into assumption we get

$$\forall \bar{y}, \bar{z}. [F_0(\bar{x}, \bar{y}) \rightarrow (\exists \bar{x}. F_0(\bar{x}, \bar{y}))] \\ \forall \bar{y}, \bar{z}. [\underbrace{(\exists \bar{x}. F_0(\bar{x}, \bar{y}))}_{H(\bar{y})} \rightarrow F_1(\bar{y}, \bar{z})]$$

$$(\forall x. \neg p(x)) \vee q$$

$$(\neg (\exists x p(x))) \vee q$$

$$(\exists x p(x)) \rightarrow q$$

and pushing  $\bar{z}$  into conclusion we get

$$\forall \bar{x}, \bar{y}. [F_0(\bar{x}, \bar{y}) \rightarrow (\forall \bar{z}. F_1(\bar{y}, \bar{z}))]$$

Given two formulas  $F_0$  and  $F_1$ , each of the formulas satisfies properties of interpolation:

- ▶  $\exists \bar{x}. F_0(\bar{x}, \bar{y})$
- ▶  $\forall \bar{z}. F_1(\bar{y}, \bar{z})$

Applying QE to them, we obtain quantifier-free interpolants.

## More on QE: One Direction to Make it More Efficient

Avoid transforming to conjunctions of literals: work directly on negation-normal form. The technique is similar to what we described for conjunctive normal form.

- + no need for DNF
- we may end up trying irrelevant bounds

This is the Cooper's algorithm:

- ▶ Reddy, Loveland: Presburger Arithmetic with Bounded Quantifier Alternation. (Gives a slight improvement of the original Cooper's algorithm.)
- ▶ Section 7.2 of the Calculus of Computation Textbook

## Eliminate Quantifiers: Example

$$\exists y. \exists x. x < -2 \wedge 1 - 5y < x \wedge 1 + y < \underline{13x}$$

$$\exists y \exists x. \quad 13x \leq -26 \wedge 13 - 65y < 13x \wedge 1 + y < 13x$$
$$z = 13x$$

$$\exists y. \exists z. z < -26 \wedge 13 - 65y < z \wedge 1 + y < z \wedge \underbrace{13 | z}_{F_1(z)}$$

$$\exists y. \bigvee_{i=1}^{13} F_1(-26-i)$$

$$\exists y. \bigvee_{i=1}^{13} (13 - 65y < -26 - i \wedge 1 + y < -26 - i \wedge 13 | -26 - i)$$

13 | -39

$$\exists y. \quad 13 - 65y < -39 \wedge 1 + y < -39$$
$$\exists y. \quad 52 < 65y \wedge y < -40$$

Check whether the formula is satisfiable

$$x < y + 2 \wedge y < x + 1 \wedge x = 3k \wedge (y = 6p + 1 \vee y = 6p - 1)$$

HW

## Apply quantifier elimination

$$\exists x. (3x + 1 < 10 \vee 7x - 6 < 7) \wedge 2 \mid x$$

AW

## Another Direction for Improvement

Handle a system of equalities more efficiently, without introducing divisibility constraints too eagerly.

Hermite normal form of an integer matrix.

$$Ax = b$$

## Eliminate variables $x$ and $y$

$$5x + 7y = a \wedge x \leq y \wedge 0 \leq x$$

HW

# Quantifier Elimination for Linear Rational Arithmetic

Consider first-order formulas with equality and  $<$  relation, interpreted over rationals.

This theory is called **dense linear order without endpoints**

For example:

$$x = |y| \quad (x = y \wedge y > 0) \vee (x = -y \wedge y \leq 0)$$

$$\forall \varepsilon. \exists \delta. (|x_1 - x_2| < \delta \wedge |y_1 - y_2| < \delta \rightarrow |3x_1 + 4y_1 - 3x_2 - 4y_2| < \varepsilon)$$

(i) Show that absolute value can be defined in first-order logic in terms of other linear operations and comparison.

# Quantifier Elimination for Linear Rational Arithmetic

Consider first-order formulas with equality and  $<$  relation, interpreted over rationals.

This theory is called **dense linear order without endpoints**

For example:

$$\forall \varepsilon. \exists \delta. (|x_1 - x_2| < \delta \wedge |y_1 - y_2| < \delta \rightarrow |3x_1 + 4y_1 - 3x_2 - 4y_2| < \varepsilon)$$

(i) Show that absolute value can be defined in first-order logic in terms of other linear operations and comparison.

Answer: replace  $F(|t|)$  with, for example

$$(t > 0 \wedge F(t)) \vee (\neg(t > 0) \wedge F(-t))$$

Is there a way to remove  $|\dots|$  while increasing formula size only linearly?

# Quantifier Elimination for Linear Rational Arithmetic

Consider first-order formulas with equality and  $<$  relation, interpreted over rationals.

This theory is called **dense linear order without endpoints**

For example:

$$\forall \varepsilon. \exists \delta. (|x_1 - x_2| < \delta \wedge |y_1 - y_2| < \delta \rightarrow |3x_1 + 4y_1 - 3x_2 - 4y_2| < \varepsilon)$$

(i) Show that absolute value can be defined in first-order logic in terms of other linear operations and comparison.

Answer: replace  $F(|t|)$  with, for example

$$(t > 0 \wedge F(t)) \vee (\neg(t > 0) \wedge F(-t))$$

Is there a way to remove  $|\dots|$  while increasing formula size only linearly?

(ii) Give quantifier elimination algorithm for this theory.

# Quantifier Elimination for Linear Rational Arithmetic

Consider first-order formulas with equality and  $<$  relation, interpreted over rationals.

This theory is called **dense linear order without endpoints**

For example:

$$\forall \varepsilon. \exists \delta. (|x_1 - x_2| < \delta \wedge |y_1 - y_2| < \delta \rightarrow |3x_1 + 4y_1 - 3x_2 - 4y_2| < \varepsilon)$$

(i) Show that absolute value can be defined in first-order logic in terms of other linear operations and comparison.

Answer: replace  $F(|t|)$  with, for example

$$(t > 0 \wedge F(t)) \vee (\neg(t > 0) \wedge F(-t))$$

Is there a way to remove  $|\dots|$  while increasing formula size only linearly?

(ii) Give quantifier elimination algorithm for this theory.

Solution is simpler than for Presburger arithmetic—no divisibility.