

Lecture 4

Paths, Triples, Postconditions, Preconditions

Viktor Kuncak

Loop-Free Programs as Relations: Summary

command c	$R(c)$	$\rho(c)$
$(x = t)$	$x' = t \wedge \bigwedge_{v \in V \setminus \{x\}} v' = v$	
$c_1 ; c_2$	$\exists \bar{z}. R(c_1)[\bar{x}' := \bar{z}] \wedge R(c_2)[\bar{x} := \bar{z}]$	$\rho(c_1) \circ \rho(c_2)$
if (*) c_1 else c_2	$R(c_1) \vee R(c_2)$	$\rho(c_1) \cup \rho(c_2)$
assume (F)	$F \wedge \bigwedge_{v \in V} v' = v$	$\Delta_{S(F)}$

$\rho(v_i = t) = \{((v_1, \dots, v_i, \dots, v_n), (v_1, \dots, v'_i, \dots, v_n)) \mid v'_i = t\}$

$S(F) = \{\bar{v} \mid F\}$, $\Delta_A = \{(\bar{v}, \bar{v}) \mid \bar{v} \in A\}$ (diagonal relation on A)

Δ (without subscript) is identity on entire set of states (no-op)

We always have: $\rho(c) = \{(\bar{v}, \bar{v}') \mid R(c)\}$

Shorthands:

$$\frac{\mathbf{if}(*)\ c_1\ \mathbf{else}\ c_2}{\mathbf{assume}(F)} \quad \Bigg| \quad c_1 \sqcap c_2$$

$$\qquad \qquad \qquad \Bigg| \quad [F]$$

Examples:

$$\mathbf{if}(F)\ c_1\ \mathbf{else}\ c_2 \equiv [F]; c_1 \sqcap [\neg F]; c_2$$

$$\mathbf{if}(F)\ c \equiv [F]; c \sqcap [\neg F]$$

Program Paths

Loop-Free Programs

c - a loop-free program whose assignments, havocs, and assumes are c_1, \dots, c_n

The relation $\rho(c)$ is of the form $E(\rho(c_1), \dots, \rho(c_n))$; it composes meanings of c_1, \dots, c_n using union (\cup) and composition (\circ)

<pre>(if (x > 0) x = x - 1 else x = 0); (if (y > 0) y = y - 1 else y = x + 1)</pre>	<pre>([x > 0]; x = x - 1 [] [¬(x>0)]; x = 0)); ([y > 0]; y = y - 1 [] [¬(y>0)]; y = x+1)</pre>	<pre>($\Delta_{S(x>0)} \circ \rho(x = x - 1)$ \cup $\Delta_{S(\neg(x>0))} \circ \rho(x = 0)$)\circ ($\Delta_{S(y>0)} \circ \rho(y = y - 1)$ \cup $\Delta_{S(\neg(y>0))} \circ \rho(y = x + 1)$)</pre>
---	--	--

Note: \circ binds stronger than \cup , so $r \circ s \cup t = (r \circ s) \cup t$

Normal Form for Loop-Free Programs

Composition distributes through union:

$$(r_1 \cup r_2) \circ (s_1 \cup s_2) = r_1 \circ s_1 \cup r_1 \circ s_2 \cup r_2 \circ s_1 \cup r_2 \circ s_2$$

Example corresponding to two if-else statements one after another:

$$\begin{aligned} & \left(\begin{array}{l} \Delta_1 \circ r_1 \\ \cup \\ \Delta_2 \circ r_2 \end{array} \right) \circ \left(\begin{array}{l} \Delta_3 \circ r_3 \\ \cup \\ \Delta_4 \circ r_4 \end{array} \right) \\ & \equiv \begin{array}{l} \Delta_1 \circ r_1 \circ \Delta_3 \circ r_3 \cup \\ \Delta_1 \circ r_1 \circ \Delta_4 \circ r_4 \cup \\ \Delta_2 \circ r_2 \circ \Delta_3 \circ r_3 \cup \\ \Delta_2 \circ r_2 \circ \Delta_4 \circ r_4 \end{array} \end{aligned}$$

Sequential composition of basic statements is called basic path.

Loop-free code describes finitely many (exponentially many) paths.

Properties of Program Contexts

Some Properties of Relations

$$(p_1 \subseteq p_2) \rightarrow (p_1 \circ p) \subseteq (p_2 \circ p)$$

$$(p_1 \subseteq p_2) \rightarrow (p \circ p_1) \subseteq (p \circ p_2)$$

$$(p_1 \subseteq p_2) \wedge (q_1 \subseteq q_2) \rightarrow (p_1 \cup q_1) \subseteq (p_2 \cup q_2)$$

$$(p_1 \cup p_2) \circ q = (p_1 \circ q) \cup (p_2 \circ q)$$

Monotonicity of Expressions using \cup and \circ

For a program with k integer variables, $S = \mathbb{Z}^k$

Consider relations that are subsets of $S \times S$ (i.e. S^2)

The set of all such relations is

$$C = \{r \mid r \subseteq S^2\}$$

Let $E(r)$ be given by any expression built from relation r and some additional relations b_1, \dots, b_n , using \cup and \circ .

Example: $E(r) = (b_1 \circ r) \cup (r \circ b_2)$

$E(r)$ is function $C \rightarrow C$, maps relations to relations

Claim: E is monotonic function on C :

$$r_1 \subseteq r_2 \rightarrow E(r_1) \subseteq E(r_2)$$

Prove or disprove.

Monotonicity of Expressions using \cup and \circ

For a program with k integer variables, $S = \mathbb{Z}^k$

Consider relations that are subsets of $S \times S$ (i.e. S^2)

The set of all such relations is

$$C = \{r \mid r \subseteq S^2\}$$

Let $E(r)$ be given by any expression built from relation r and some additional relations b_1, \dots, b_n , using \cup and \circ .

Example: $E(r) = (b_1 \circ r) \cup (r \circ b_2)$

$E(r)$ is function $C \rightarrow C$, maps relations to relations

Claim: E is monotonic function on C :

$$r_1 \subseteq r_2 \rightarrow E(r_1) \subseteq E(r_2)$$

Prove or disprove.

Proof: induction on the expression tree defining E , using monotonicity properties of \cup and \circ

Union-Distributivity of Expressions using \cup and \circ

Claim: E distributes over unions, that is, if $r_i, i \in I$ is a family of relations,

$$E\left(\bigcup_{i \in I} r_i\right) = \bigcup_{i \in I} E(r_i)$$

Prove or disprove.

Union-Distributivity of Expressions using \cup and \circ

Claim: E distributes over unions, that is, if $r_i, i \in I$ is a family of relations,

$$E\left(\bigcup_{i \in I} r_i\right) = \bigcup_{i \in I} E(r_i)$$

Prove or disprove.

False. Take $E(r) = r \circ r$ and consider relations r_1, r_2 . The claim becomes

$$(r_1 \cup r_2) \circ (r_1 \cup r_2) = r_1 \circ r_1 \cup r_2 \circ r_2$$

that is,

$$r_1 \circ r_1 \cup r_1 \circ r_2 \cup r_2 \circ r_1 \cup r_2 \circ r_2 = r_1 \circ r_1 \cup r_2 \circ r_2$$

Taking, for example, $r_1 = \{(1, 2)\}$, $r_2 = \{(2, 3)\}$ we obtain

$$\{(1, 3)\} = \emptyset \quad (\text{false})$$

Union “Distributivity” in One Direction

Lemma:

$$E\left(\bigcup_{i \in I} r_i\right) \supseteq \bigcup_{i \in I} E(r_i)$$

Union “Distributivity” in One Direction

Lemma:

$$E\left(\bigcup_{i \in I} r_i\right) \supseteq \bigcup_{i \in I} E(r_i)$$

Proof. Let $r = \bigcup_{i \in I} r_i$. Note that, for every i , $r_i \subseteq r$. We have shown that E is monotonic, so $E(r_i) \subseteq E(r)$. Since all $E(r_i)$ are included in $E(r)$, so is their union, so

$$\bigcup_{i \in I} E(r_i) \subseteq E(r)$$

as desired.

Union-Distributivity - Refined

Does distributivity

$$E\left(\bigcup_{i \in I} r_i\right) = \bigcup_{i \in I} E(r_i)$$

hold, for each of these cases

1. If $E(r)$ is given by an expression containing r at most once?

Union-Distributivity - Refined

Does distributivity

$$E\left(\bigcup_{i \in I} r_i\right) = \bigcup_{i \in I} E(r_i)$$

hold, for each of these cases

1. If $E(r)$ is given by an expression containing r at most once?
Proof: Induction on expression for $E(r)$. Only one branch of the tree may contain r . Note previous counter-example uses r twice.

Union-Distributivity - Refined

Does distributivity

$$E\left(\bigcup_{i \in I} r_i\right) = \bigcup_{i \in I} E(r_i)$$

hold, for each of these cases

1. If $E(r)$ is given by an expression containing r at most once?
Proof: Induction on expression for $E(r)$. Only one branch of the tree may contain r . Note previous counter-example uses r twice.
2. If $E(r)$ contains r any number of times, but I is a set of natural numbers and r_i is an increasing sequence:
 $r_1 \subseteq r_2 \subseteq r_3 \subseteq \dots$

Union-Distributivity - Refined

Does distributivity

$$E\left(\bigcup_{i \in I} r_i\right) = \bigcup_{i \in I} E(r_i)$$

hold, for each of these cases

1. If $E(r)$ is given by an expression containing r at most once?
Proof: Induction on expression for $E(r)$. Only one branch of the tree may contain r . Note previous counter-example uses r twice.
2. If $E(r)$ contains r any number of times, but I is a set of natural numbers and r_i is an increasing sequence:
 $r_1 \subseteq r_2 \subseteq r_3 \subseteq \dots$. Induction. In the previous counter-example the largest relation will contain all other $r_i \circ r_j$.
3. If $E(r)$ contains r any number of times, but $r_i, i \in I$ is a **directed family** of relations: for each i, j there exists k such that $r_i \cup r_j \subseteq r_k$, and I is possibly uncountably infinite.

Union-Distributivity - Refined

Does distributivity

$$E\left(\bigcup_{i \in I} r_i\right) = \bigcup_{i \in I} E(r_i)$$

hold, for each of these cases

1. If $E(r)$ is given by an expression containing r at most once?
Proof: Induction on expression for $E(r)$. Only one branch of the tree may contain r . Note previous counter-example uses r twice.
2. If $E(r)$ contains r any number of times, but I is a set of natural numbers and r_i is an increasing sequence:
 $r_1 \subseteq r_2 \subseteq r_3 \subseteq \dots$. Induction. In the previous counter-example the largest relation will contain all other $r_i \circ r_j$.
3. If $E(r)$ contains r any number of times, but $r_i, i \in I$ is a **directed family** of relations: for each i, j there exists k such that $r_i \cup r_j \subseteq r_k$, and I is possibly uncountably infinite.
Induction. Generalizes the previous case.

About Strength and Weakness

Putting Conditions on Sets Makes them Smaller

Let P_1 and P_2 be formulas (“conditions”) whose free variables are among \bar{x} . Those variables may denote program state.

When we say “condition P_1 is stronger than condition P_2 ” it simply means

$$\forall \bar{x}. (P_1 \rightarrow P_2)$$

- ▶ if we know P_1 , we immediately get (conclude) P_2
- ▶ if we know P_2 we need not be able to conclude P_1

Stronger condition = smaller set: if P_1 is stronger than P_2 then

$$\{\bar{x} \mid P_1\} \subseteq \{\bar{x} \mid P_2\}$$

- ▶ strongest possible condition: “false” \rightsquigarrow smallest set: \emptyset
- ▶ weakest condition: “true” \rightsquigarrow biggest set: set of all tuples

Hoare Triples

About Hoare Logic


We have seen how to translate programs into relations. We will use these relations in a proof system called Hoare logic. Hoare logic is a way of inserting annotations into code to make proofs about (imperative) program behavior simpler.

Example proof:

```
//{0 <= y}
i = y;
//{0 <= y & i = y}
r = 0;
//{0 <= y & i = y & r = 0}
while //{r = (y-i)*x & 0 <= i}
  (i > 0) (
    //{r = (y-i)*x & 0 < i}
    r = r + x;
    //{r = (y-i+1)*x & 0 < i}
    i = i - 1
    //{r = (y-i)*x & 0 <= i}
  )
//{r = x * y}
```

Hoare Triple and Friends

Sir Charles Antony Richard Hoare



Sir Charles Antony Richard Hoare giving a conference at the EPFL on 20 June 2011

Born 11 January 1934

$$P, Q \subseteq S \quad r \subseteq S \times S$$

Hoare Triple:

$$\{P\} r \{Q\} \iff \forall s, s' \in S. (s \in P \wedge (s, s') \in r \rightarrow s' \in Q)$$

$\{P\}$ does not denote a singleton set containing P but is just a notation for an “assertion” around a command. Likewise for $\{Q\}$.

Strongest postcondition:

$$sp(P, r) = \{s' \mid \exists s. s \in P \wedge (s, s') \in r\}$$

Weakest precondition:

$$wp(r, Q) = \{s \mid \forall s'. (s, s') \in r \rightarrow s' \in Q\}$$

Exercise: Which Hoare triples are valid?

Assume all variables to be over integers.

1. $\{j = a\} j := j+1 \{a = j + 1\}$
2. $\{i = j\} i := j+i \{i > j\}$
3. $\{j = a + b\} i := b; j := a \{j = 2 * a\}$
4. $\{i > j\} j := i+1; i := j+1 \{i > j\}$
5. $\{i \neq j\} \text{ if } i > j \text{ then } m := i - j \text{ else } m := j - i \{m > 0\}$
6. $\{i = 3*j\} \text{ if } i > j \text{ then } m := i - j \text{ else } m := j - i \{m - 2*j = 0\}$

Postconditions and Their Strength

What is the relationship between these postconditions?

$$\{x = 5\} \quad x := x + 2 \quad \{\mathbf{x} > \mathbf{0}\}$$

$$\{x = 5\} \quad x := x + 2 \quad \{\mathbf{x} = \mathbf{7}\}$$

Postconditions and Their Strength

What is the relationship between these postconditions?

$$\{x = 5\} \quad x := x + 2 \quad \{x > 0\}$$

$$\{x = 5\} \quad x := x + 2 \quad \{x = 7\}$$

- ▶ weakest conditions (predicates) correspond to largest sets
- ▶ strongest conditions (predicates) correspond to smallest sets

that satisfy a given property.

(Graphically, a stronger condition $x > 0 \wedge y > 0$ denotes one quadrant in plane, whereas a weaker condition $x > 0$ denotes the entire half-plane.)


Strongest Postconditions

Strongest Postcondition

Definition: For $P \subseteq S$, $r \subseteq S \times S$,

$$sp(P, r) = \{s' \mid \exists s. s \in P \wedge (s, s') \in r\}$$

This is simply the relation image of a set.




Weakest Preconditions

Weakest Precondition

Definition: for $Q \subseteq S$, $r \subseteq S \times S$,

$$wp(r, Q) = \{s \mid \forall s'. (s, s') \in r \rightarrow s' \in Q\}$$

Note that this is in general not the same as $sp(Q, r^{-1})$ when the relation is non-deterministic or partial.



Three Forms of Hoare Triple

Lemma: the following three conditions are equivalent:

- ▶ $\{P\}r\{Q\}$
- ▶ $P \subseteq wp(r, Q)$
- ▶ $sp(P, r) \subseteq Q$

Three Forms of Hoare Triple

Lemma: the following three conditions are equivalent:

- ▶ $\{P\}r\{Q\}$
- ▶ $P \subseteq wp(r, Q)$
- ▶ $sp(P, r) \subseteq Q$

Proof. The three conditions expand into the following three formulas


- ▶ $\forall s, s'. [(s \in P \wedge (s, s') \in r) \rightarrow s' \in Q]$
- ▶ $\forall s. [s \in P \rightarrow (\forall s'. (s, s') \in r \rightarrow s' \in Q)]$
- ▶ $\forall s'. [(\exists s. s \in P \wedge (s, s') \in P) \rightarrow s' \in Q]$

which are easy to show equivalent using basic first-order logic properties.

Lemma: Characterization of sp

$sp(P, r)$ is the the smallest set Q such that $\{P\}r\{Q\}$, that is:

- ▶ $\{P\}r\{sp(P, r)\}$
- ▶ $\forall Q \subseteq S. \{P\}r\{Q\} \rightarrow sp(P, r) \subseteq Q$



$$\{P\} r \{Q\} \Leftrightarrow \forall s, s' \in S. (s \in P \wedge (s, s') \in r \rightarrow s' \in Q)$$

$$sp(P, r) = \{s' \mid \exists s. s \in P \wedge (s, s') \in r\}$$

Proof of Lemma: Characterization of sp


Apply Three Forms of Hoare triple. The two conditions then reduce to:

- ▶ $sp(P, r) \subseteq sp(P, r)$
- ▶ $\forall P \subseteq S. sp(P, r) \subseteq Q \rightarrow sp(P, r) \subseteq Q$

Lemma: Characterization of wp

$wp(r, Q)$ is the largest set P such that $\{P\}r\{Q\}$, that is:

- ▶ $\{wp(r, Q)\}r\{Q\}$
- ▶ $\forall P \subseteq S. \{P\}r\{Q\} \rightarrow P \subseteq wp(r, Q)$



$$\{P\}r\{Q\} \Leftrightarrow \forall s, s' \in S. (s \in P \wedge (s, s') \in r \rightarrow s' \in Q)$$

$$wp(r, Q) = \{s \mid \forall s'. (s, s') \in r \rightarrow s' \in Q\}$$

Proof of Lemma: Characterization of wp

Apply Three Forms of Hoare triple. The two conditions then reduce to:

- ▶ $wp(r, Q) \subseteq wp(r, Q)$
- ▶ $\forall P \subseteq S. P \subseteq wp(r, Q) \rightarrow P \subseteq wp(r, Q)$

Exercise: Postcondition of inverse versus wp

Lemma:

$$S \setminus wp(r, Q) = sp(S \setminus Q, r^{-1})$$

In other words, when instead of good states we look at the complement set of “error states”, then wp corresponds to doing sp backwards.

Note that $r^{-1} = \{(y, x) \mid (x, y) \in r\}$ and is always defined.

Exercise: Postcondition of inverse versus wp

Lemma:

$$S \setminus wp(r, Q) = sp(S \setminus Q, r^{-1})$$

In other words, when instead of good states we look at the complement set of “error states”, then wp corresponds to doing sp backwards.

Note that $r^{-1} = \{(y, x) \mid (x, y) \in r\}$ and is always defined.

Proof of the lemma: Expand both sides and apply basic first-order logic properties.

More Laws on Preconditions and Postconditions

Disjunctivity of sp

$$sp(P_1 \cup P_2, r) = sp(P_1, r) \cup sp(P_2, r)$$

$$sp(P, r_1 \cup r_2) = sp(P, r_1) \cup sp(P, r_2)$$

Conjunctivity of wp

$$wp(r, Q_1 \cap Q_2) = wp(r, Q_1) \cap wp(r, Q_2)$$

$$wp(r_1 \cup r_2, Q) = wp(r_1, Q) \cap wp(r_2, Q)$$

Pointwise wp

$$wp(r, Q) = \{s \mid s \in S \wedge sp(\{s\}, r) \subseteq Q\}$$

Pointwise sp

$$sp(P, r) = \bigcup_{s \in P} sp(\{s\}, r)$$

Hoare Logic for Loop-free Code

Expanding Paths

The condition

$$\{P\} \left(\bigcup_{i \in J} r_i \right) \{Q\}$$

is equivalent to

$$\forall i. i \in J \rightarrow \{P\} r_i \{Q\}$$

Proof: By definition, or use that the first condition is equivalent to $sp(P, \bigcup_{i \in J} r_i) \subseteq Q$ and $\{P\} r_i \{Q\}$ to $sp(P, r_i) \subseteq Q$

Transitivity

If $\{P\} s_1 \{Q\}$ and $\{Q\} s_2 \{R\}$ then also $\{P\} s_1 \circ s_2 \{R\}$.

We write this as the following inference rule:

$$\frac{\{P\} s_1 \{Q\}, \quad \{Q\} s_2 \{R\}}{\{P\} s_1 \circ s_2 \{R\}}$$

Exercise

We call a relation $r \subseteq S \times S$ functional if

$\forall x, y, z \in S. (x, y) \in r \wedge (x, z) \in r \rightarrow y = z$. For each of the following statements either give a counterexample or prove it. In the following, $Q \subseteq S$.

- (i) for any r , $wp(r, S \setminus Q) = S \setminus wp(r, Q)$
- (ii) if r is functional, $wp(r, S \setminus Q) = S \setminus wp(r, Q)$
- (iii) for any r , $wp(r, Q) = sp(Q, r^{-1})$
- (iv) if r is functional, $wp(r, Q) = sp(Q, r^{-1})$
- (v) for any r , $wp(r, Q_1 \cup Q_2) = wp(r, Q_1) \cup wp(r, Q_2)$
- (vi) if r is functional, $wp(r, Q_1 \cup Q_2) = wp(r, Q_1) \cup wp(r, Q_2)$
- (vii) for any r , $wp(r_1 \cup r_2, Q) = wp(r_1, Q) \cup wp(r_2, Q)$
- (viii) Alice has a conjecture: For all sets S and relations $r \subseteq S \times S$ it holds:

$$\left(S \neq \emptyset \wedge \text{dom}(r) = S \wedge \Delta_S \cap r = \emptyset \right) \rightarrow \left(r \circ r \cap ((S \times S) \setminus r) \neq \emptyset \right)$$

where $\Delta_S = \{(x, x) \mid x \in S\}$, $\text{dom}(r) = \{x \mid \exists y. (x, y) \in r\}$. She tried many sets and relations and did not find any counterexample. Is her conjecture true? If so, prove it; if false, provide a counterexample for which S is as small as possible.